

GREEDY BASES FOR BESOV SPACES

S. J. DILWORTH, D. FREEMAN, E. ODELL, AND TH. SCHLUMPRECHT

ABSTRACT. We prove that the Banach space $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$, which is isomorphic to certain Besov spaces, has a greedy basis whenever $1 \leq p \leq \infty$ and $1 < q < \infty$. Furthermore, the Banach spaces $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_1}$, with $1 < p \leq \infty$, and $(\oplus_{n=1}^{\infty} \ell_p^n)_{c_0}$, with $1 \leq p < \infty$ do not have a greedy bases. We prove as well that the space $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$ has a 1-greedy basis if and only if $1 \leq p = q \leq \infty$.

1. INTRODUCTION

Let X be a Banach space and let (x_i) be a Schauder basis for X with biorthogonal sequence (x_i^*) . For $x \in X$ and $n \geq 1$, the error in the best n -term approximation to x (using (x_i)) is given by

$$\sigma_n(x) := \inf \left\{ \left\| x - \sum_{i \in A} a_i x_i \right\| : (a_i) \subset \mathbb{R}, |A| \leq n \right\}.$$

Let $A_n(x) \subset \mathbb{N}$ be the indices corresponding to a choice of n largest coefficients of x in absolute value, i.e. $A_n(x)$ satisfies

$$\min \{ |e_i^*(x)| : i \in A_n(x) \} \geq \max \{ |e_i^*(x)| : i \in \mathbb{N} \setminus A_n(x) \}.$$

Then $G_n(x) := \sum_{i \in A_n(x)} x_i^*(x) x_i$ is called an n^{th} *greedy approximant* to x . We say that (x_i) is *greedy* with constant C if

$$\|x - G_n(x)\| \leq C \sigma_n(x) \quad (x \in X, n \geq 1).$$

If $C = 1$ then (x_i) is said to be 1-greedy. Temlyakov [T1] proved that the Haar system for $L_p[0, 1]^d$ ($1 < p < \infty$, $d \geq 1$) is greedy, which provides an important theoretical justification for the thresholding procedure used in data compression. Subsequently, Konyagin and Temlyakov [KT] gave a very useful abstract characterization of greedy bases. To state their result, we recall that (x_i) is *unconditional* with constant K if, for all choices of signs, we have

$$\left\| \sum_{i=1}^{\infty} \pm x_i^*(x) x_i \right\| \leq K \|x\| \quad (x \in X).$$

Key words and phrases. Greedy bases; Besov spaces.

2000 Mathematics Subject Classification: Primary 46B15, Secondary 41A65.

The research of the first, third, and fourth authors was supported by National Science Foundation grants DMS 0701552, DMS 0700126, and DMS 0856148, respectively. The second author was supported by grant N000140811113 of the Office of Naval Research. The first and third authors were also supported by the Workshop in Analysis and Probability at Texas A&M University in 2009.

We say that (x_i) is *democratic* with constant Δ if, for all finite $A, B \subset \mathbb{N}$ with $|A| = |B|$, we have

$$\left\| \sum_{i \in A} x_i \right\| \leq \Delta \left\| \sum_{i \in B} x_i \right\|.$$

Theorem A. [KT] *Suppose that (x_i) is unconditional with constant K and democratic with constant Δ . Then (x_i) is greedy with constant $K + K^3\Delta$. Conversely, if (x_i) is greedy with constant C then (x_i) is unconditional with constant C and democratic with constant C^2 .*

Theorem A was used in [W1, W2] to prove that $L_p[0, 1]$ ($p \neq 2$) has a greedy basis that is not equivalent to a subsequence of the Haar basis, and in [DHK] to prove that ℓ_p and $L_p[0, 1]$ ($p \neq 2$) have a continuum of mutually non-equivalent greedy bases. It was also used in [DKKT] to study duality for greedy bases, and a similar theorem was proved in [DKKT] to characterize the larger class of *almost greedy* bases (see also [DKK]).

Some examples of greedy bases are given in [W2]. In most cases these bases are greedy simply because they are *symmetric* (e.g. Riesz bases for a Hilbert space, which are equivalent to the unit vector basis of ℓ_2 , or good wavelet bases for the Besov spaces $B_{\alpha,p}^p(\mathbb{R})$, which are equivalent to the unit vector basis of ℓ_p), or because they are equivalent to the Haar basis (e.g. good wavelet bases for $L_p(\mathbb{R}^d)$) or to a subsequence of the Haar basis (e.g. generalized Haar systems [K]). In [GH] certain wavelet bases in the Triebel-Lizorkin spaces $f_{p,r}^s$ are shown to be greedy. In [AW] it is proved that 1-symmetric bases (e.g. the unit vector bases of Orlicz and Lorentz sequence spaces) are in fact 1-greedy. On the other hand, there are examples of spaces with an unconditional basis but no democratic unconditional basis, and hence no greedy basis, e.g. certain spaces with a unique unconditional basis up to permutation [BCLT], the spaces $\ell_p \oplus \ell_q$ and $\ell_p \oplus c_0$ for $1 \leq p < q < \infty$ [AW], and the original Tsirelson space T^* [T2]. Wojtaszczyk [W3] proved that the L_p spaces ($1 < p < \infty$) are the only rearrangement-invariant function spaces on $[0, 1]$ for which the Haar system is greedy.

Using Theorem A we prove that for every $1 \leq p \leq \infty$ and $1 < q < \infty$ the Banach space $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$ has a greedy basis. Furthermore, we show that the Banach space $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_1}$ does not have a greedy basis whenever $1 < p \leq \infty$. This answers a question posed by P. Wojtaszczyk, who asked when such spaces have a greedy basis. The problem of finding a greedy basis for Banach spaces of the form $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$ is particularly pertinent from the approximation theoretical standpoint as such spaces are isomorphic to certain Besov spaces on the circle [R]. As shown in [R, Theorem 2] (see also [Pi, page 255]), the Besov space $B_{p,q}^{\alpha,m}[0, 1]$, where $1 \leq p \leq \infty$, $1 \leq q < \infty$, $m \in \{-1, 0, 1, 2, \dots\}$, and $\alpha \in (1, m + 1 + 1/p)$, is isomorphic to $\ell_1^{m+2} \oplus (\oplus_{n=0}^{\infty} \ell_p^{2^n})_{\ell_q}$ which is easily seen to be isomorphic to $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$.

The greedy bases which we construct in Theorem 1 differ from the examples discussed above in that they are neither subsymmetric (see [LT, p. 114] for this notion) nor equivalent to a subsequence of the Haar basis.

The following result completely characterizes for which pairs (p, q) the space $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$ has a greedy basis.

Theorem 1. *Let $1 \leq p, q \leq \infty$.*

- a) *If $1 < q < \infty$ then the Banach space $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$ has a greedy basis.*

- b) The spaces $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_1}$, with $1 < p \leq \infty$, and $(\oplus_{n=1}^{\infty} \ell_p^n)_{c_0}$, with $1 \leq p < \infty$, do not have greedy bases.

The following result yields that only in the trivial case that $p = q$ does $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$ have a 1-greedy basis.

Theorem 2. *Let $1 \leq p \leq \infty$ and let $(E_n)_{n=1}^{\infty}$ be a sequence of finite dimensional Banach spaces. If $(x_i)_{i=1}^{\infty}$ is a normalized 1-greedy basis for the space $(\oplus_{n=1}^{\infty} E_n)_{\ell_p}$ then $(x_i)_{i=1}^{\infty}$ is 1-equivalent to the standard unit vector basis for ℓ_p (as usual, if $p = \infty$ we consider the c_0 -sum).*

As the cases $\ell_p \oplus \ell_q$ and $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$ are settled the following spaces might be interesting to consider.

Problem 3. Assume $1 < p \neq q < \infty$. Does $\ell_q(\ell_p) = (\oplus \ell_p)_{\ell_q}$ have a greedy basis?

We thank P. Wojtaszczyk for his comments on a preliminary version of our paper and for drawing [R] to our notice.

2. PROOF OF THEOREMS 1 AND 2

Part (a) of Theorem 1 will follow easily from the following Lemma, whose proof will require some work.

Lemma 4. *Let $1 \leq p \leq \infty$ and $1 < q < \infty$ and let $\varepsilon > 0$. There is a constant $1 \leq K < \infty$ such that for all $N \in \mathbb{N}$ there exist $M = M_N$ and a finite normalized sequence $(x_i)_{i=1}^M \subset \ell_q(\ell_p^N)$ such that*

- a) $(x_i)_{i=1}^M$ is 1-unconditional,
- b) $(1 - \varepsilon)|A| \leq \left\| \sum_{i \in A} x_i \right\|^q \leq (1 + \varepsilon)|A|$ for all $A \subset \{1, \dots, M\}$,
- c) the span of $(x_i)_{i=1}^M$ is K -complemented in $\ell_q(\ell_p^N)$, and
- d) ℓ_p^N is isometric to a K -complemented subspace of the span of $(x_i)_{i=1}^M$.

Using the lemma, we give a quick proof of the first part of Theorem 1.

Proof of Theorem 1 (a). Let $1 \leq p \leq \infty$ and $1 < q < \infty$. It will be more convenient for us to work with the space $X := (\oplus_{N=1}^{\infty} \ell_q(\ell_p^N))_{\ell_q}$ instead of $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$. That these spaces are isomorphic follows from *Pelczyński's Decomposition Method* [Pe], which says that if two Banach spaces are complementably embedded in each other, and one of them is isomorphic to the (countably infinite) ℓ_r -sum, $1 \leq r < \infty$, or c_0 -sum of itself, then they are isomorphic. It is easy to observe that X and $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$ are 1-complemented in each other, and that X is isometric to its ℓ_q sum.

We let $\varepsilon > 0$ and choose, for each $N \in \mathbb{N}$, a sequence $(x_i^{(N)})_{i=1}^{M_N}$ in the N th coordinate of $X = (\oplus_{N=1}^{\infty} \ell_q(\ell_p^N))_{\ell_q}$ which satisfies Lemma 4. From the conditions (a) and (b) in Lemma 4 it follows that $(x_i^{(N)})_{N \in \mathbb{N}, 1 \leq i \leq M_N}$ is a 1-unconditional and $\frac{1+\varepsilon}{1-\varepsilon}$ -democratic sequence in X . As the span of $(x_i^{(N)})_{i=1}^{M_N}$ is K -complemented in $\ell_q(\ell_p^N)$, for each $N \in \mathbb{N}$, it follows that the closed span of $(x_i^{(N)})_{N \in \mathbb{N}, 1 \leq i \leq M_N}$ is K -complemented in X . Furthermore, as ℓ_p^N is K -complemented in the span of $(x_i^{(N)})_{i=1}^{M_N}$ for each $N \in \mathbb{N}$, it follows that $(\oplus_{N=1}^{\infty} \ell_p^N)_{\ell_q}$ is K -complemented in

the closed span of $(x_i^{(N)})_{N \in \mathbb{N}, 1 \leq i \leq M_N}$. Thus by the Pełczyński decomposition theorem, X is isomorphic to the closed span of $(x_i^{(N)})_{N \in \mathbb{N}, 1 \leq i \leq M_N}$. Hence X , and thus $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_q}$, has a greedy basis. $\square$

Proof of Lemma 4. For $\varepsilon > 0$, we choose numbers $\varepsilon_i \searrow 0$ such that $\prod_{i=1}^{\infty} (1 + \varepsilon_i) < 1 + \varepsilon$ and $\prod_{i=1}^{\infty} (1 - \varepsilon_i) > 1 - \varepsilon$. For each $n \in \mathbb{N}$, we denote by $(e_{(i,n)})_{i=1}^N$ the unit vector basis for the n th coordinate of $\ell_q(\ell_p^N)$, and we denote by $(e_{(i,n)}^*)_{i=1}^N$ their biorthogonal functionals. Thus the norm on $\ell_q(\ell_p^N)$ is calculated by

$$\left\| \sum a_{(i,n)} e_{(i,n)} \right\| = \left(\sum_{n=1}^{\infty} \left(\sum_{i=1}^N |a_{(i,n)}|^p \right)^{q/p} \right)^{1/q} \text{ for } (a_{(i,n)} : 1 \leq i \leq N, n \in \mathbb{N}) \subset \mathbb{R}.$$

If $x \in \ell_q(\ell_p^N)$, then we denote the support of x by $\text{supp}(x) = \{(i, n) | e_{(i,n)}^*(x) \neq 0\}$.

Before proceeding we fix two sequences of integers $(m_i), (k_i) \in [\mathbb{N}]^{\omega}$ with $m_1 = k_1 = 1$ and which satisfy the following inequalities for all $i > 1$.

- i) $1/\varepsilon_i < m_i$,
- ii) $m_i^{1/q} + 1 < (1 + \varepsilon_i)^{1/q} m_i^{1/q}$,
- iii) $(1 + (m_i/k_i)^{1/q})^q < 1 + \varepsilon_i$, and
- iv) $1 - \varepsilon_i < (1 - (m_i/k_i)^{1/q})^q$.

The above inequalities can be easily guaranteed by first choosing m_i large enough to satisfy i) and ii), and then choosing k_i large enough to satisfy iii) and iv). For the sake of convenience we define $n_j = \prod_{i=1}^j k_i$ for all $j \in \mathbb{N}$. We define the finite family $(x_{(i,j)})_{1 \leq i \leq N, 1 \leq j \leq n_N/n_i}$ by

$$x_{(i,j)} = \frac{1}{n_i^{1/q}} \sum_{s=1}^{n_i} e_{(i,s+(j-1)n_i)} \quad \text{for all } 1 \leq i \leq N \text{ and } 1 \leq j \leq n_N/n_i.$$

It is clear that $(x_{(i,j)})$ is a normalized and 1-unconditional basic sequence as the sequence has pairwise disjoint support. Also, $\bigcup_{i \leq N, j \leq n_N/n_i} \text{supp}(x_{(i,j)}) = \{1, 2, \dots, N\} \times \{1, 2, \dots, n_N\}$. For each integer $1 \leq \ell \leq N$ and subset

$$A \subset \{(i, j) \in \mathbb{N}^2 | 1 \leq i \leq \ell \text{ and } 1 \leq j \leq n_N/n_i\},$$

we will prove by induction on ℓ that

$$(1) \quad |A| \prod_{i=2}^{\ell} (1 - \varepsilon_i) \leq \left\| \sum_{(i,j) \in A} x_{(i,j)} \right\|^q \leq |A| \prod_{i=2}^{\ell} (1 + \varepsilon_i).$$

First note that if $\ell = 1$ and $A \subset \{(1, j) \in \mathbb{N}^2 | 1 \leq j \leq n_N\}$ then

$$\left\| \sum_{(1,j) \in A} x_{(1,j)} \right\|^q = \left\| \sum_{(1,j) \in A} e_{(1,j)} \right\|^q = |A|.$$

Thus (1) is trivially satisfied. We now assume that equation (1) is satisfied for a given $1 \leq \ell < N$ and we will prove it for $\ell + 1$. We first partition the set $\Omega = \{(i, j) \in \mathbb{N}^2 : 1 \leq i \leq \ell + 1 \text{ and } 1 \leq j \leq n_N/n_i\}$, into sets $\Omega_1, \Omega_2, \dots, \Omega_{n_N/n_{\ell+1}}$ defined for each $1 \leq r \leq n_N/n_{\ell+1}$ by

$$\Omega_r := \{(i, j) \in \mathbb{N}^2 | 1 \leq i \leq \ell + 1, (r - 1)n_{\ell+1}/n_i + 1 \leq j \leq rn_{\ell+1}/n_i\}.$$

Observe that for $1 \leq r \leq n_N/n_{\ell+1}$, $1 \leq i \leq \ell + 1$, and $1 \leq j \leq n_N/n_i$

$$\begin{aligned} (i, j) \in \Omega_n &\iff (r-1)\frac{n_{\ell+1}}{n_i} + 1 \leq j \leq r\frac{n_{\ell+1}}{n_i} \\ &\iff \text{supp}(x_{(i,j)}) = \{i\} \times [(j-1)n_i + 1, jn_i] \subset \{i\} \times [(r-1)n_{\ell+1} + 1, rn_{\ell+1}]. \end{aligned}$$

Since $\text{supp}(x_{(\ell+1,r)}) = \{\ell+1\} \times [(r-1)n_{\ell+1} + 1, rn_{\ell+1}]$, it follows that

$$(i, j) \in \Omega_r \iff \{(\ell+1, s) : (i, s) \in \text{supp}(x_{(i,j)})\} \subset \text{supp}(x_{(\ell+1,r)}).$$

Given the set $A \subset \Omega$, we partition A into $A_1, A_2, \dots, A_{n_N/n_{\ell+1}}$, by defining $A_r = A \cap \Omega_r$, for all $1 \leq r \leq n_N/n_{\ell+1}$. We note that

$$\left\| \sum_{(i,j) \in A} x_{(i,j)} \right\|^q = \sum_{r=1}^{n_N/n_{\ell+1}} \left\| \sum_{(i,j) \in A_r} x_{(i,j)} \right\|^q$$

and that $(x_{(i,j)})_{(i,j) \in \Omega_1}$ is 1-equivalent to $(x_{(i,j)})_{(i,j) \in \Omega_r}$, for all $1 \leq r \leq n_N/n_{\ell+1}$. Thus, to prove the inequality (1), we just need to consider the case $A = A_1$. We first note that if $(\ell+1, 1) \notin A_1$, then the inequality (1) is immediately true by the induction hypothesis. Thus we now assume that $(\ell+1, 1) \in A_1$ and $A_1 \setminus \{(\ell+1, 1)\} \neq \emptyset$.

Roughly speaking, we will argue that either $|A_1|$ is large enough so that $\sum_{(i,j) \in A_1} x_{(i,j)}$ can be replaced by $\sum_{(i,j) \in A_1 \setminus \{(\ell+1, 1)\}} x_{(i,j)}$, or $|A_1|$ is so small that a large part of the support of $x_{(\ell+1, 1)}$ is disjoint from

$$B_1 = \{(\ell+1, n) : (i, n) \in \text{supp}(x_{(i,j)}) \text{ for some } j \in \{1, 2, \dots, n_N/n_i\}\},$$

and we can approximate $x_{(\ell+1, 1)}$ by its projection onto $\text{span}(e_{(\ell+1, n)} : n \in \{1, 2, \dots, N\} \setminus B_1)$.

The first case we consider is that $|A_1| \geq m_{\ell+1}$. This assumption, together with the inequality $m_{\ell+1} > 1/\varepsilon_{\ell+1}$, yields

$$\frac{|A_1|}{|A_1| - 1} \leq \frac{m_{\ell+1}}{m_{\ell+1} - 1} < \frac{1}{1 - \varepsilon_{\ell+1}}.$$

This allows us to obtain the desired lower estimate. Indeed,

$$\left\| \sum_{(i,j) \in A_1} x_{(i,j)} \right\| \geq \left\| \sum_{(i,j) \in A_1 \setminus \{(\ell+1, 1)\}} x_{(i,j)} \right\| \geq (|A_1| - 1) \prod_{i=2}^{\ell} (1 - \varepsilon_i) \geq |A_1| \prod_{i=2}^{\ell+1} (1 - \varepsilon_i).$$

To prove the upper estimate in (1), we use that $|A_1| \geq m_{\ell+1}$ together with $ii)$ to get

$$\frac{(|A_1| - 1)^{1/q} + 1}{|A_1|^{1/q}} < \frac{m_{\ell+1}^{1/q} + 1}{m_{\ell+1}^{1/q}} < (1 + \varepsilon_{\ell+1})^{1/q}.$$

Thus,

$$\begin{aligned} \left\| \sum_{(i,j) \in A_1} x_{(i,j)} \right\| &\leq \left\| \sum_{(i,j) \in A_1 \setminus \{(\ell+1, 1)\}} x_{(i,j)} \right\| + 1 \\ &\leq (|A_1| - 1)^{1/q} \left(\prod_{i=2}^{\ell} (1 + \varepsilon_i) \right)^{1/q} + \left(\prod_{i=2}^{\ell} (1 + \varepsilon_i) \right)^{1/q} < |A_1|^{1/q} \left(\prod_{i=2}^{\ell+1} (1 + \varepsilon_i) \right)^{1/q}. \end{aligned}$$

This completes the proof of (1) for $\ell + 1$ in the case that $|A_1| \geq m_{\ell+1}$. We now assume that $|A_1| < m_{\ell+1}$. The size of the support of each $x_{(i,j)}$ is given by $|\text{supp}(x_{(i,j)})| = n_i$ for all $1 \leq i \leq \ell + 1$ and $1 \leq j \leq n_N/n_i$. We thus have a simple estimate for the size of the union of the supports,

$$(2) \quad |\cup_{(i,j) \in A_1 \setminus \{(\ell+1,1)\}} \text{supp}(x_{(i,j)})| = \sum_{(i,j) \in A_1 \setminus \{(\ell+1,1)\}} |\text{supp}(x_{(i,j)})| < |A_1| n_\ell.$$

We define sets

$$B_1 := \{(\ell + 1, n) \in \mathbb{N}^2 : (m, n) \in \cup_{(i,j) \in A_1 \setminus \{(\ell+1,1)\}} \text{supp}(x_{(i,j)}) \text{ for some } 1 \leq m \leq \ell\} \text{ and} \\ B_2 := \text{supp}(x_{(\ell+1,1)}) \setminus B_1.$$

The inequality (2) gives that $|B_1| < |A_1| n_\ell$. For $i = 1, 2$, we define $P_{B_i} x_{(\ell+1,1)}$ by $P_{B_i} x_{(\ell+1,1)} = \frac{1}{n_{\ell+1}^{1/q}} \sum_{(\ell+1,j) \in B_i} e_{(\ell+1,j)}$. We may estimate the value $\|P_{B_1} x_{(\ell+1,1)}\|$ by

$$(3) \quad \|P_{B_1} x_{(\ell+1,1)}\| = \frac{1}{n_{\ell+1}^{1/q}} |B_1|^{1/q} < \left(\frac{|A_1| n_\ell}{n_{\ell+1}} \right)^{1/q} < \left(\frac{m_{\ell+1} n_\ell}{n_{\ell+1}} \right)^{1/q} = \left(\frac{m_{\ell+1}}{k_{\ell+1}} \right)^{1/q}.$$

We use this to obtain the following estimate.

$$\begin{aligned} \left\| \sum_{(i,j) \in A_1} x_{(i,j)} \right\|^q &= \left\| P_{B_2} x_{(\ell+1,1)} + P_{B_1} x_{(\ell+1,1)} + \sum_{(i,j) \in A_1 \setminus \{(\ell+1,1)\}} x_{(i,j)} \right\|^q \\ &= \|P_{B_2} x_{(\ell+1,1)}\|^q + \left\| P_{B_1} x_{(\ell+1,1)} + \sum_{(i,j) \in A_1 \setminus \{(\ell+1,1)\}} x_{(i,j)} \right\|^q \\ &\leq 1 + \left[\left(\frac{m_{\ell+1}}{k_{\ell+1}} \right)^{1/q} + \left((|A_1| - 1) \prod_{i=1}^{\ell} (1 + \varepsilon_i) \right)^{1/q} \right]^q \\ &= 1 + \left[1 + \left(\frac{m_{\ell+1}}{k_{\ell+1}} \right)^{1/q} \left((|A_1| - 1) \prod_{i=1}^{\ell} (1 + \varepsilon_i) \right)^{-1/q} \right]^q (|A_1| - 1) \prod_{i=1}^{\ell} (1 + \varepsilon_i) \\ &\leq \left[1 + \left(\frac{m_{\ell+1}}{k_{\ell+1}} \right)^{1/q} \left(\prod_{i=1}^{\ell} (1 + \varepsilon_i) \right)^{-1/q} \right]^q \prod_{i=1}^{\ell} (1 + \varepsilon_i) \\ &\quad + \left[1 + \left(\frac{m_{\ell+1}}{k_{\ell+1}} \right)^{1/q} \left((|A_1| - 1) \prod_{i=1}^{\ell} (1 + \varepsilon_i) \right)^{-1/q} \right]^q (|A_1| - 1) \prod_{i=1}^{\ell} (1 + \varepsilon_i) \\ &\leq \left[1 + \left(\frac{m_{\ell+1}}{k_{\ell+1}} \right)^{1/q} \left(\prod_{i=1}^{\ell} (1 + \varepsilon_i) \right)^{-1/q} \right]^q |A_1| \prod_{i=1}^{\ell} (1 + \varepsilon_i) \\ &\leq |A_1| \prod_{i=1}^{\ell+1} (1 + \varepsilon_i) \quad \text{by iii).} \end{aligned}$$

For proving the remaining lower inequality in (1), we will use the following estimate for $\|P_{B_2}x_{(\ell+1,1)}\|$ which follows from (3).

$$\|P_{B_2}x_{(\ell+1,1)}\| \geq 1 - \|P_{B_1}x_{(\ell+1,1)}\| > 1 - \left(\frac{m_{\ell+1}}{k_{\ell+1}}\right)^{1/q}.$$

This yields

$$\begin{aligned} \left\| \sum_{(i,j) \in A_1} x_{(i,j)} \right\|^q &= \|P_{B_2}x_{(\ell+1,1)}\|^q + \left\| P_{B_1}x_{(\ell+1,1)} + \sum_{(i,j) \in A_1 \setminus \{(\ell+1,1)\}} x_{(i,j)} \right\|^q \\ &\geq \|P_{B_2}x_{(\ell+1,1)}\|^q + \left(-\|P_{B_1}x_{(\ell+1,1)}\| + \left\| \sum_{(i,j) \in A_1 \setminus \{(\ell+1,1)\}} x_{(i,j)} \right\| \right)^q \\ &= \|P_{B_2}x_{(\ell+1,1)}\|^q \\ &\quad + \left(1 - \|P_{B_1}x_{(\ell+1,1)}\| \cdot \left\| \sum_{(i,j) \in A_1 \setminus \{(\ell+1,1)\}} x_{(i,j)} \right\|^{-1} \right)^q \left\| \sum_{(i,j) \in A_1 \setminus \{(\ell+1,1)\}} x_{(i,j)} \right\|^q \\ &\geq \left(1 - \left(\frac{m_{\ell+1}}{k_{\ell+1}}\right)^{1/q} \right)^q + \left(1 - \left(\frac{m_{\ell+1}}{k_{\ell+1}}\right)^{1/q} \right)^q (|A_1| - 1) \prod_{i=1}^{\ell} (1 - \varepsilon_i) \\ &\geq \left(1 - \left(\frac{m_{\ell+1}}{k_{\ell+1}}\right)^{1/q} \right)^q |A_1| \prod_{i=1}^{\ell} (1 - \varepsilon_i) \\ &> \prod_{i=1}^{\ell+1} (1 - \varepsilon_i) |A_1| \quad \text{by iv).} \end{aligned}$$

Thus we have proven the inequalities (1) in all cases. It remains to prove that there exists a constant $1 \leq K < \infty$, independent of $N \in \mathbb{N}$, such that $X := \text{span}(x_{(i,j)})$ is K -complemented in $\ell_q(\ell_p^N)$ and ℓ_p^N is isometric to a K -complemented subspace of X . For each $1 \leq i \leq N$, we define the vector y_i as

$$y_i := \frac{n_i^{1/q}}{n_N^{1/q}} \sum_{j=1}^{n_N/n_i} x_{(i,j)} = \frac{1}{n_N^{1/q}} \sum_{j=1}^{n_N} e_{(i,j)}.$$

It should be clear that $(y_i)_{i=1}^N$ is 1-equivalent to the unit vector basis for ℓ_p^N . Indeed, if $(a_i) \in \ell_p^N$ then

$$\left\| \sum_{i=1}^N a_i y_i \right\| = \left(\sum_{j=1}^{n_N} \left(\sum_{i=1}^N \frac{|a_i|^p}{n_N^{p/q}} \right)^{q/p} \right)^{1/q} = \left(\frac{1}{n_N} \sum_{j=1}^{n_N} \left(\sum_{i=1}^N |a_i|^p \right)^{q/p} \right)^{1/q} = \left(\sum_{i=1}^N |a_i|^p \right)^{1/p}.$$

We let $Y = \text{span}(y_i)$ and define projections $T_X : \ell_q(\ell_p^N) \rightarrow X$ and $T_Y : \ell_q(\ell_p^N) \rightarrow Y$ by

$$T_X \left(\sum a_{(i,j)} e_{(i,j)} \right) = \sum_{i=1}^N \sum_{j=1}^{n_N/n_i} \sum_{s=1}^{n_i} \left(\frac{1}{n_i} \sum_{k=1}^{n_i} a_{(i,k+(j-1)n_i)} \right) e_{(i,s+(j-1)n_i)}$$

$$\begin{aligned}
&= \sum_{i=1}^N \sum_{j=1}^{n_N/n_i} \left(\frac{1}{n_i^{(q-1)/q}} \sum_{k=1}^{n_i} a_{(i,k+(j-1)n_i)} \right) x_{(i,j)} \text{ and} \\
T_Y \left(\sum a_{(i,j)} e_{(i,j)} \right) &= \sum_{i=1}^N \sum_{s=1}^{n_N} \left(\frac{1}{n_N} \sum_{k=1}^{n_N} a_{(i,k)} \right) e_{(i,s)} = \sum_{i=1}^N \left(\frac{1}{n_N^{(q-1)/q}} \sum_{k=1}^{n_N} a_{(i,k)} \right) y_i.
\end{aligned}$$

It is simple to check that T_X and T_Y are projections of $\ell_q(\ell_p^N)$ onto X and Y respectively. As Y is a subspace of X , we have that T_Y restricted to X is a projection of X onto Y . Thus we just need to prove that there exists a uniform constant K such that $\|T_X\|, \|T_Y\| \leq K$. We note that $T_X = T_Y$ if, considering a more general case, $n_i = n_N$ for all $1 \leq i \leq N$. Thus proving there exists a constant K independent of $(n_i)_{i=1}^N$ and N such that $\|T_X\| \leq K$ will prove the inequality for $\|T_Y\|$ as well. We first consider the case that $p = q$. In this case, the basis $(e_{(i,j)})$ for the space $\ell_q(\ell_q^N)$ is symmetric. The operator T_X is then an averaging operator on a space with a symmetric basis, and hence has norm one.

Now if $q < p < \infty$, then $\ell_q(\ell_p^N)$ is an interpolation space for the spaces $\ell_q(\ell_q^N)$ and $\ell_q(\ell_\infty^N)$. Then by the vector valued Riesz-Thorin interpolation theorem [BP] (see also [C]) , if $\theta = \frac{q}{p}$ then

$$\|T_X\|_{\ell_q(\ell_p^N)} \leq \|T_X\|_{\ell_q(\ell_q^N)}^\theta \|T_X\|_{\ell_q(\ell_\infty^N)}^{1-\theta} \leq \|T_X\|_{\ell_q(\ell_\infty^N)}.$$

Thus if we prove that there exists a uniform constant K such that $\|T_X\|_{\ell_q(\ell_\infty^N)} \leq K$ then the result will follow as well for all $1 < q < p < \infty$. On the other hand, if $1 \leq p < q < \infty$ then $1 < q' < p' \leq \infty$, with $q' = q/(q-1)$ and $p' = p/(p-1)$. It is simple to check that our operator $T_X : \ell_q(\ell_p^N) \rightarrow \ell_q(\ell_p^N)$ has adjoint $T_X^* = T_X : \ell_{q'}(\ell_{p'}^N) \rightarrow \ell_{q'}(\ell_{p'}^N)$. We thus have that $\|T_X\|_{\ell_q(\ell_p^N)} = \|T_X^*\|_{\ell_{q'}(\ell_{p'}^N)} \leq K$.

All that remains is to prove is that there exists a uniform constant K such that $\|T_X\|_{\ell_q(\ell_\infty^N)} \leq K$. This constant K will come from a *discretization* of the classical Hardy-Littlewood maximal operator [HL], which is defined as

$$(4) \quad n_u(g)(x) = \sup_{y < x < z} \frac{1}{z-y} \int_y^z |g(t)| dt \quad \text{for } x \in \mathbb{R} \text{ and } g \in L_{loc}^1(\mathbb{R}).$$

It is known that the operator n_u is of strong type (q, q) for $1 < q \leq \infty$ and for a proof of this see [G, Theorem 8.9.1 and Corollary 8.9.1]. In other words, there exists a constant $1 \leq K < \infty$ such that

$$\left(\int |n_u(g)(x)|^q dx \right)^{1/q} \leq K \left(\int |g(x)|^q dx \right)^{1/q} \text{ for all } g \in L_q(\mathbb{R}).$$

By applying this to step functions whose discontinuities are contained in $\mathbb{N}$, we get the following inequality for ℓ_q .

$$(5) \quad \left(\sum_{j \in \mathbb{N}} \left(\sup_{m \leq j \leq n} \frac{1}{n-m+1} \sum_{k=m}^n |a_k| \right)^q \right)^{1/q} \leq K \left(\sum_{j \in \mathbb{N}} |a_j|^q \right)^{1/q} \text{ for all } (a_j) \in \ell_q.$$

We now prove that $\|T_X\|_{\ell_q(\ell_\infty^N)} \leq K$. As $\ell_q(\ell_\infty^N)$ is reflexive, the operator T_X attains it's norm at an extreme point of the ball $B_{\ell_q(\ell_\infty^N)}$. The set of extreme points of $B_{\ell_q(\ell_\infty^N)}$ is given

by

$$\text{Ext}(B_{\ell_q(\ell_\infty^N)}) = \left\{ \sum_{j=1}^{\infty} a_j \sum_{i=1}^N \varepsilon_{(i,j)} e_{(i,j)} : \varepsilon_{(i,j)} = \pm 1, (a_j) \in S_{\ell_q} \right\}.$$

Thus there exists constants $\varepsilon_{(i,j)} = \pm 1$ and a sequence $(a_j) \in S_{\ell_q}$ such that

$$\begin{aligned} \|T_X\|_{\ell_q(\ell_\infty^N)} &= \left\| T_X \left(\sum_{j=1}^{\infty} a_j \sum_{i=1}^N \varepsilon_{(i,j)} e_{(i,j)} \right) \right\| \\ &= \left\| \sum_{i=1}^N \sum_{j=1}^{n_N/n_i} \sum_{s=1}^{n_i} \left(\frac{1}{n_i} \sum_{k=1}^{n_i} \varepsilon_{(i,k+(j-1)n_i)} a_{k+(j-1)n_i} \right) e_{(i,s+(j-1)n_i)} \right\| \\ &\leq \left\| \sum_{i=1}^N \sum_{j=1}^{n_N/n_i} \sum_{s=1}^{n_i} \left(\frac{1}{n_i} \sum_{k=1}^{n_i} |a_{k+(j-1)n_i}| \right) e_{(i,s+(j-1)n_i)} \right\| \\ &\leq \left\| \sum_{i=1}^N \sum_{j=1}^{n_N/n_i} \sum_{s=1}^{n_i} \left(\sup_{m \leq 1+(j-1)n_i \leq n} \frac{1}{n+1-m} \sum_{k=m}^n |a_k| \right) e_{(i,s+(j-1)n_i)} \right\| \\ &\leq \left\| \sum_{i=1}^N \sum_{j=1}^{n_N/n_i} \left(\sup_{m \leq j \leq n} \frac{1}{n+1-m} \sum_{k=m}^n |a_k| \right) e_{(i,j)} \right\| \\ &= \left(\sum_{j=1}^{n_N} \left(\sup_{m \leq j \leq n} \frac{1}{n+1-m} \sum_{k=m}^n |a_k| \right)^q \right)^{1/q} \\ &\leq K \left(\sum_{j=1}^{n_N} |a_k|^q \right)^{1/q} \quad \text{by (5).} \end{aligned}$$

□

Theorem 5. *The Banach space $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_1}$ does not have a greedy basis whenever $1 < p \leq \infty$.*

We recall that Bourgain, Casazza, Lindenstrauss, and Tzafriri proved that the spaces $(\oplus_{n=1}^{\infty} \ell_2^n)_{\ell_1}$ and $(\oplus_{n=1}^{\infty} \ell_\infty^n)_{\ell_1}$ each have unconditional bases which are unique up to permutation [BCLT]. In particular, these spaces cannot have a greedy basis, as they each have an unconditional basis which is not greedy. Thus we need only to prove Theorem 5 for the case $1 < p < \infty$. This is important for us as ℓ_p has non-trivial type and cotype when $1 < p < \infty$. We rely on the following proposition, which was used in [BCLT] to prove, among other uniqueness results, that $(\oplus_{n=1}^{\infty} \ell_2^n)_{\ell_1}$ has a unique unconditional basis up to permutation.

Proposition B. [BCLT, Proposition 2.1] *Let V be a Banach lattice of type p and cotype q , for some $1 \leq p \leq q < \infty$ and let $C_c, C_u \geq 1$ be constants. There exists a uniform constant $K \geq 1$ which satisfies the following statement. If Z is a C_c -complemented subspace of the direct sum $X = (\oplus_{n=1}^{\infty} V)_{c_0}$ and Z has a normalized basis $(z_n)_{n=1}^k$ with unconditional constant C_u , then there exists a partition of the integers $\{1, 2, \dots, k\}$ into mutually disjoint*

subsets $\{\tau_s\}_{s=1}^r$ so that, for any choice of scalars $\{\alpha_n\}_{n=1}^k$, we have

$$K^{-1} \max_{1 \leq s \leq r} \left(\sum_{n \in \tau_s} |\alpha_n|^q \right)^{1/q} \leq \left\| \sum_{n=1}^k \alpha_n z_n \right\| \leq K \max_{1 \leq s \leq r} \left(\sum_{n \in \tau_s} |\alpha_n|^p \right)^{1/p}.$$

Using Proposition B, we are now prepared to give a proof of Theorem 5.

Proof of Theorem 5. Let $1 < p < \infty$. To reach a contradiction, we assume that $X = (\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_1}$ has a normalized basis $(x_i)_{i=1}^{\infty}$ which is C_d -democratic and C_u -unconditional for some constants $C_d, C_u \geq 1$. Let $(x_i^*) \subset (\oplus_n \ell_p^n)^*_{\ell_1} = (\oplus_n \ell_q^n)_{\ell_{\infty}}$, with $\frac{1}{p} + \frac{1}{q} = 1$, be the biorthogonal functionals. We let $(e_{(i,n)})_{1 \leq i \leq n < \infty}$ to be the unit vector basis for X with biorthogonal functionals $(e_{(i,n)}^*)_{1 \leq i \leq n < \infty}$. By a standard perturbation argument we may assume that

$$\text{supp}(x_j) = \{(i, n) : n \in \mathbb{N}, i \leq n, e_{(i,n)}^*(x_j) \neq 0\}$$

is finite for each $j \in \mathbb{N}$.

We now fix $N \in \mathbb{N}$. There exists $M_N \in \mathbb{N}$ such that $(x_j)_{j=1}^N \subset (\oplus_{n=1}^{M_N} \ell_p^n)_{\ell_1}$, i.e. $e_{(i,n)}^*(x_j) = 0$ for all $n > M_N$ and $i \leq n$. Define $(z_{(N,j)}^*)_{j=1}^N \subset (\oplus_{n=1}^{M_N} \ell_p^n)^*_{\ell_1}$ by $z_{(N,j)}^* = x_j^*|_{(\oplus_{n=1}^{M_N} \ell_p^n)_{\ell_1}}$ for all $1 \leq j \leq N$. As $(x_j)_{j=1}^{\infty}$ has basis constant at most C_u , it is easy to show both that $(x_j^*)_{j=1}^N$ is C_u -equivalent to $(z_{(N,j)}^*)_{j=1}^N$, and that the span of $(z_{(N,j)}^*)_{j=1}^N$ is C_u -complemented in $(\oplus_{n=1}^{M_N} \ell_p^n)^*_{\ell_1}$. Indeed, let $J : \text{span}(x_i^*)_{i=1}^N \rightarrow X$ be the inclusion map. Then $J^* : X^* \rightarrow \text{span}(x_i^*)_{i=1}^N$ is a quotient map of norm 1. Let $H : \text{span}(x_i^*)_{i=1}^N \rightarrow \text{span}(z_{(N,i)}^*)_{i=1}^N$ be the isomorphism defined by $H(x_i^*) = z_{(N,i)}^*$, which has norm at most C_u . Then $H \circ J^* : X^* \rightarrow \text{span}(z_{(N,i)}^*)_{i=1}^N$ is a quotient map of norm at most C_u . We just need to check that $H \circ J^*(z_{(N,i)}^*) = z_{(N,i)}^*$. Indeed, for $x \in \text{span}(x_j : j \leq N)$, $J^*(z_{(N,i)}^*)(x) = z_{(N,i)}^*(x) = x_i^*(x)$. Hence, $J^*(z_{(N,i)}^*) = x_i^*$ and $H \circ J^*(z_{(N,i)}^*) = z_{(N,i)}^*$. Thus $H \circ J^*$ is a projection of norm at most C_u .

We will be applying Proposition B for the space $V = \ell_q$ and consider the spaces $\text{span}(z_{(N,i)}^* : i \leq N)$, $N \in \mathbb{N}$, (in the natural way) to be C_u -complemented subspaces of $c_0(\ell_q)$. For the sake of convenience, we denote $\underline{q} = \min(q, 2)$ and $\bar{q} = \max(q, 2)$. We have thus defined $\underline{q}$ and $\bar{q}$ exactly so that ℓ_q has type $\underline{q}$ and cotype $\bar{q}$. By Proposition B, there exists a constant K independent of $N \in \mathbb{N}$, and there exists for all $N \in \mathbb{N}$ a partition of the integers $\{1, 2, \dots, N\}$ into mutually disjoint subsets $\{\tau_s^N\}_{s=1}^{r_N}$ so that, for any choice of scalars $\{\alpha_n\}_{n=1}^N$, we have

$$(6) \quad K^{-1} \max_{1 \leq s \leq r_N} \left(\sum_{n \in \tau_s^N} |\alpha_n|^{\bar{q}} \right)^{1/\bar{q}} \leq \left\| \sum_{n=1}^N \alpha_n z_{(N,n)}^* \right\| \leq K \max_{1 \leq s \leq r_N} \left(\sum_{n \in \tau_s^N} |\alpha_n|^{\underline{q}} \right)^{1/\underline{q}}.$$

We first consider the case that $\sup_{N \in \mathbb{N}} \max_{1 \leq s \leq r_N} |\tau_s^N| < \infty$. In this case we have that if $N \in \mathbb{N}$ and $(\alpha_i)_{i=1}^N \subset \mathbb{R}$ then

$$\left\| \sum_{i=1}^N \alpha_i x_i^* \right\| \leq C_u \left\| \sum_{i=1}^N \alpha_i z_{(N,i)}^* \right\| \leq C_u K \left(\sup_{M \in \mathbb{N}} \max_{1 \leq s \leq r_M} |\tau_s^M| \right) \max_{1 \leq i \leq N} |\alpha_i|.$$

Hence (x_i^*) is equivalent to the unit vector basis of c_0 , which implies that (x_i) is equivalent to the unit vector basis of ℓ_1 . This is a contradiction as (x_i) is a basis for $X = (\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_1}$, and X is not isomorphic to ℓ_1 as $1 < p < \infty$. We now consider the remaining case that $\sup_{N \in \mathbb{N}} \max_{1 \leq s \leq r_N} |\tau_s^N| = \infty$.

First note that there exists a subsequence of $(x_i)_{i=1}^{\infty}$ which is equivalent to the unit vector basis of ℓ_1 . Indeed, there is a subsequence (x'_j) which converges in the w^* -topology (considering X as the dual of $(\oplus_{n=1}^{\infty} \ell_q^n)_{c_0}$) to some x . If $x \neq 0$, then it follows from the assumed unconditionality that (x'_j) is equivalent to the unit vector basis of ℓ_1 , and if $x = 0$, then we can find a perturbation (z'_j) of a further subsequence, so that for each $n \in \mathbb{N}$, there is at most one j so that the ℓ_q^n component of z'_j is not 0. But this also implies that there is a subsequence equivalent to the unit vector basis of ℓ_1 .

Thus as (x_i) is democratic, there is a $C \geq 1$ so that $C \left\| \sum_{i \in A} x_i \right\| > |A|$ for all finite sets $A \subset \mathbb{N}$. We choose $N \in \mathbb{N}$ and $1 \leq s \leq r_N$ such that $|\tau_s^N|^{1/\bar{q}} > 2KCC_u^2$. We have the following estimates.

$$\begin{aligned} \left\| \sum_{i \in \tau_s^N} x_i \right\| &\leq 2C_u \left(\sum_{i \in \tau_s^N} b_i x_i^* \right) \left(\sum_{i \in \tau_s^N} x_i \right) \quad \text{for some } (b_i) \in c_{00}, \text{ with } \left\| \sum_{i \in \tau_s^N} b_i x_i^* \right\| = 1 \\ &\leq 2C_u^2 \frac{\sum_{i \in \tau_s^N} |b_i|}{\left\| \sum_{i \in \tau_s^N} b_i z_{(N,i)}^* \right\|} \\ &\leq 2KC_u^2 \frac{\sum_{i \in \tau_s^N} |b_i|}{\sum_{i \in \tau_s^N} |b_i|^{\bar{q}})^{1/\bar{q}}} \\ &\leq 2KC_u^2 |\tau_s^N|^{1-1/\bar{q}} \quad \text{by Hölder's inequality.} \end{aligned}$$

Combining this result with the inequality $|\tau_s^N|^{1/\bar{q}} > 2KCC_u^2$ gives the following contradiction.

$$2KCC_u^2 |\tau_s^N|^{1-1/\bar{q}} < |\tau_s^N| < C \left\| \sum_{i \in \tau_s^N} x_i \right\| \leq 2KCC_u^2 |\tau_s^N|^{1-1/\bar{q}}$$

As both possible cases result in a contradiction, we see that $(\oplus_{n=1}^{\infty} \ell_p^n)_{\ell_1}$ cannot have a greedy basis when $1 < p < \infty$. $\square$

Finally the case $q = \infty$ and $1 \leq p < \infty$ in Theorem 1 is easy to handle.

Proposition 6. *For $1 \leq p < \infty$ the space $(\oplus \ell_p^n)_{c_0}$ does not have a greedy basis.*

Proof. Assume that (x_n) is a greedy basis of $(\oplus \ell_p^n)_{c_0}$. Since (x_n) is democratic and must contain a subsequence which is equivalent to the unit vector basis of c_0 it follows that for some constant $C \geq 1$ that,

$$\left\| \sum_{n \in A} x_n \right\| \leq C, \text{ for all finite } A \subset \mathbb{N}.$$

This together with the unconditionality of (x_n) implies that (x_n) is equivalent to the unit basis of c_0 , which is a contradiction since $(\oplus \ell_p^n)_{c_0}$ is not isomorphic to c_0 . $\square$

We have now finished the proof of Theorem 1 and determined which spaces of the form $(\oplus_{N=1}^{\infty} \ell_p^N)_{\ell_q}$ have a greedy basis. We now turn to the proof of Theorem 2.

We rely on the concept of greedy permutations developed by Albiac and Wojtaszczyk [AW], which we recall here. Let $M(x)$ denote the subset of the support of x consisting of the largest coordinates of x in absolute value. We will say that a one to one map $\pi : \text{supp}(x) \rightarrow \mathbb{N}$ is a greedy permutation of x if $\pi(j) = j$ for all $j \in \text{supp}(x) \setminus M(x)$ and if $j \in M(x)$ then, either $\pi(j) = j$ or $\pi(j) \notin \text{supp}(x)$.

Definition. A basic sequence (e_n) is defined to have property (A) if for any $x \in \text{span}(e_i)$ we have

$$\left\| \sum_{n \in \text{supp}(x)} e_n^*(x) e_n \right\| = \left\| \sum_{n \in \text{supp}(x)} \theta_{\pi(n)} e_n^*(x) e_{\pi(n)} \right\|$$

for all greedy permutations π of x and all sequences of signs (θ_k) with $\theta_{\pi(n)} = 1$ if $\pi(n) = n$.

We recall that (e_n) is called *C-suppression unconditional*, for some $C \geq 1$, if for any $(a_i) \subset c_{00}$ and any $A \subset \mathbb{N}$,

$$\left\| \sum_{i \in A} a_i e_i \right\| \leq C \left\| \sum_{i \in \mathbb{N}} a_i e_i \right\|.$$

Theorem C. [AW, Theorem 3.4] *A basic sequence (e_n) is 1-greedy if and only if (e_n) is 1-suppression unconditional and satisfies property (A).*

Proof of Theorem 2. We first consider the case that $1 < p < \infty$. We show that if $A \subset \mathbb{N}$ is any finite set, then $\left\| \sum_{i \in A} a_i x_i \right\| = (\sum_{i \in A} |a_i|^p)^{1/p}$ for all $(a_i) \in c_{00}$. This is trivial if $|A| = 1$ as (x_i) is normalized. We now assume that the equality holds for $|A| \leq k$ for some $k \geq 1$. Let $(a_i) \in c_{00}$ and $A \subset \mathbb{N}$ such that $|A| = k + 1$. Choose $N \in A$ such that $|a_N| = \max_{i \in A} |a_i|$. We define $\pi_j : A \rightarrow \mathbb{N}$ by $\pi_j(N) = j$ and $\pi_j(n) = n$ for all $n \neq N$. The map π_j is a greedy permutation whenever $j \notin A$, and hence by Theorem C we have the following equalities.

$$\begin{aligned} \left\| \sum_{i \in A} a_i x_i \right\| &= \left\| \sum_{i \in A, i \neq N} a_i x_i + a_N x_j \right\| \quad \text{for all } j \notin A \\ &= \lim_{j \rightarrow \infty} \left\| \sum_{i \in A, i \neq N} a_i x_i + a_N x_j \right\| \\ &= \left(\left\| \sum_{i \in A, i \neq N} a_i x_i \right\|^p + |a_N|^p \right)^{1/p}, \quad \text{as } (x_j) \text{ is normalized and weakly null,} \\ &= \left(\sum_{i \in A} |a_i|^p \right)^{1/p}, \quad \text{by the induction hypothesis.} \end{aligned}$$

This finishes the proof of the induction step, and, thus, the proof of our claim.

The case $p = \infty$, in which case we consider the c_0 -sum of the E_n , works similarly, as every normalized unconditional sequence in $(\sum E_n)_{c_0}$ must be weakly null.

We now consider the ℓ_1 case. Let (x_i) be a 1-greedy basis for $(\sum E_n)_{\ell_1}$. If (x_i) is w^* -null with respect to the w^* topology given by $(\sum E_n^*)_{\ell_\infty}$, then the proof that (x_i) is 1-equivalent

to the unit vector basis for ℓ_1 is the same as the previous case $1 < p < \infty$. If (x_i) is not w^* -null, then (x_i) has a subsequence (x_{k_i}) which converges w^* to some non-zero $x \in (\sum E_n)_{\ell_1}$. Hence $(x_{k_i} - x)$ is w^* -null. This implies that $\lim_{i \rightarrow \infty} \|y + x_{k_i} - x\| = \|y\| + \lim_{i \rightarrow \infty} \|x_{k_i} - x\|$ for all $y \in (\sum E_n)_{\ell_1}$. We use this to achieve the following equalities.

$$\begin{aligned} \lim_{n \rightarrow \infty} \lim_{i \rightarrow \infty} \|x_{k_n} - x_{k_i}\| &= \lim_{n \rightarrow \infty} \lim_{i \rightarrow \infty} \|(x_{k_n} - x) - (x_{k_i} - x)\| \\ &= \lim_{n \rightarrow \infty} \|x_{k_n} - x\| + \lim_{i \rightarrow \infty} \|x_{k_i} - x\| = 2 \lim_{i \rightarrow \infty} \|x_{k_i} - x\|. \end{aligned}$$

Furthermore,

$$\begin{aligned} \lim_{n \rightarrow \infty} \lim_{i \rightarrow \infty} \|x_{k_n} + x_{k_i}\| &= \lim_{n \rightarrow \infty} \lim_{i \rightarrow \infty} \|(x_{k_n} - x) + (x_{k_i} - x) + 2x\| \\ &= \lim_{n \rightarrow \infty} \|x_{k_n} - x + 2x\| + \lim_{i \rightarrow \infty} \|x_{k_i} - x\| \\ &= 2\|x\| + \lim_{n \rightarrow \infty} \|x_{k_n} - x\| + \lim_{i \rightarrow \infty} \|x_{k_i} - x\| = 2\|x\| + 2 \lim_{i \rightarrow \infty} \|x_{k_i} - x\|. \end{aligned}$$

As (x_i) is 1-greedy, we must have, by Theorem C, that $\lim_{n \rightarrow \infty} \lim_{i \rightarrow \infty} \|x_{k_n} - x_{k_i}\| = \lim_{n \rightarrow \infty} \lim_{i \rightarrow \infty} \|x_{k_n} + x_{k_i}\|$. This however implies that $\|x\| = 0$, which is a contradiction with our assumption that x is non-zero. $\square$

REFERENCES

- [AW] F. Albiac and P. Wojtaszczyk, *Characterization of 1-greedy bases*, J. Approx. Theory, **138** no.1 (2006), 65-86.
- [BCLT] J. Bourgain, P.G. Casazza, J. Lindenstrauss, and L. Tzafriri, *Banach spaces with a unique unconditional basis, up to permutation*, Memoirs Amer. Math. Soc. No. 322, 1985.
- [BP] A. Benedek and R. Panzone, *The space L_p , with mixed norm*. Duke Math. J. **28** (1961) 301-324.
- [C] A. P. Calderón, *Intermediate spaces, and Interpolation*, Studia Math. **24** (1964), 113-190.
- [DHK] S. J. Dilworth, Mark Hoffmann and Denka Kutzarova *Non-equivalent greedy and almost-greedy bases in ℓ_p* , J. Funct. Spaces and Appl., **318** (2006), 692-706.
- [DKK] S. J. Dilworth, N. J. Kalton and Denka Kutzarova, *On the existence of almost greedy bases in Banach spaces*, Studia Math. (Pełczyński Anniversary Volume), Studia Math. **159** (2003), 67-101.
- [DKKT] S. J. Dilworth, N. J. Kalton, Denka Kutzarova, and V. N. Temlyakov, *The Thresholding Greedy Algorithm, Greedy Bases, and Duality*, Constr. Approx. **19** (2003), no. 4, 575-595.
- [EW] I. S. Edelstein and P. Wojtaszczyk, *On projections and unconditional bases in direct sums of Banach spaces*, Studia Math. **56** (1976), no. 3, 263-276.
- [G] D.J.H. Garling, *Inequalities: a journey into linear analysis* Cambridge University Press, 2007.
- [GH] Gustavo Garrigós and Eugenio Hernández, *Sharp Jackson and Bernstein Inequalities for N -Term Approximation in Sequence Spaces with Applications*, Indiana Univ. Math. J. **53** (2004), 1739-1762.
- [HL] G. H. Hardy and J. E. Littlewood, *A maximal theorem with function-theoretic applications*, Acta Math. **54** (1930), 81-116.
- [K] A. Kamont, *General Haar systems and greedy approximation*, Studia Math. **145** (2001), 165-184.
- [KT] S.V. Konyagin and V.N. Temlyakov, *A remark on greedy approximation in Banach spaces*, East J. Approx., **5** no.3 (1999), 365-379.
- [LT] J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces I: Sequence spaces*, Springer-Verlag, Berlin-Heidelberg-New York, 1977.
- [Pe] A. Pełczyński, *Projections in certain Banach spaces*, Studia Math. **19** (1960), 209 - 228.
- [Pi] A. Pietsch, *Eigenvalues and s -numbers*. Cambridge Studies in Advanced Mathematics, 13. Cambridge University Press, Cambridge, 1987.

- [R] S. Ropela, *Spline bases in Besov spaces*, Bull. Acad. Pol. Sci., Ser. Sci. Math. Astron. Phys., **24** (1976), 319–325.
- [T1] V. N. Temlyakov, *The best m -term approximation and greedy algorithms*, Adv. Comput. Math. **8** (1998), 249–265.
- [T2] B. S. Tsirelson, *Not every Banach space contains ℓ_p or c_0* , Functional Anal. Appl. **8** (1974), 138–141.
- [W1] P. Wojtaszczyk, *Greedy algorithm for general biorthogonal systems*, J. Approx. Theory **107** (2000), 293–314.
- [W2] P. Wojtaszczyk, *Greedy type bases in Banach spaces*, Constructive theory of functions, 136–155, DARBA, Sofia, 2003.
- [W3] P. Wojtaszczyk, *Greediness of the Haar system in rearrangement invariant spaces*. Approximation and probability, 385–395, Banach Center Publ., 72, Polish Acad. Sci., Warsaw, 2006.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SC 29208, USA
E-mail address: `dilworth@math.sc.edu`

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843, USA
E-mail address: `freeman@math.tamu.edu`

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS, 1 UNIVERSITY STATION C1200, AUSTIN, TX 78712, USA
E-mail address: `odell@math.utexas.edu`

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843, USA
E-mail address: `thomas.schlumprecht@math.tamu.edu`